



Microneedling with PRP and/or PRP LIFT FILLER

Description of the procedure

This treatment involves the collection of your blood (approximately 8-16ml). After blood is collected it is spun down using a centrifuge to separate out the plasma and platelet portion using the separator gel as a special filter. The PRP portion of our blood is then used at the point of care to re-energize your cells into rejuvenating themselves. The product is 100% your own blood by-product (autologous). If you have any questions please do not hesitate to ask your provider.

EXCLUSIONS, PRECAUTIONS AND WARNINGS OF TREATMENT

- Autoimmune therapies
- Keloid scarring
- History of eczema, psoriasis and other chronic conditions
- Patients with active herpes, acne or rosacea
- History of actinic keratosis, raised moles or warts on targeted area
- History of hemophilia, irregular blood pressure
- Susceptibility to capillary ectasia due to steroid use for extended periods
- Scleroderma
- Collagen vascular disease
- Cardiac abnormalities, pacemakers, blood thinners
- Active bacterial or fungal infection
- Immunosuppression
- Facial melanosis
- Malignant tumors
- History of any type of cancer or suspicious lesions or moles in treatment areas.

- Pregnant or nursing
- Recent Botox, collagen, fat or other methods of augmentation with injected materials

PROCEDURE

- The procedure is typically completed in 30-60 minutes.
- The treatment time can vary based on the required treatment and anatomical site being treated.
- Numbing cream will be applied for 15-20 minutes prior to treatment
- The level of pain and discomfort varies with a person's tolerance and both may be experienced during treatment with gradual cessation of pain after treatment. .
- Localized pinpoint bleeding into the treatment area can occur and will resolve within 24 hours.
- Immediately following the procedure the most commonly reported temporary side effects are redness, swelling, bruising, tenderness, tingling, numbness, lumpiness, and/or feelings of pressure or fullness at the injection site and/or in the treated area.
- To avoid bruising, Avoid alcohol consumption for a minimum of 6 hours and refrain from taking blood thinners such as Aspirin and NSAIDs for several days. Tylenol is recommended if needed for discomfort.

AFTER PROCEDURE CARE

- Avoid applying lotions, creams, perfumes, powders or makeup to the treated area for at least 24 hours post treatment.
- Avoid touching or scrubbing at the injection sites for 24 hours after treatment.
- To maximize results for a minimum of 5 hours after PRP has been applied to your skin, do not wash your skin, expose the treated area to direct high heat or engage in activities that will get the skin wet or cause you to sweat.
- Avoid hot baths or showers, massages, rigorous exercise and/or any activity requiring direct skin contact or causing excessive perspiration in the treatment area 24-48hrs after treatment.
- No tanning or sun exposure.
- Wear sunscreen SPF 30 or higher with reapplying every 2 hours for 24 hours after treatment.
- A skin routine will be suggested and individualized depending on patient's needs by provider.